

Discrete realization of group symmetric LOCC-detection of maximally entangled state

Masahito Hayashi^{1,*}

¹*Graduate School of Information Sciences, Tohoku University, Aoba-ku, Sendai, 980-8579, Japan*

Group symmetric LOCC measurement for detecting maximally entangled state is considered. Usually, this type measurement has continuous-valued outcomes. However, any realizable measurement has finite-valued outcomes. This paper proposes discrete realizations of such a group symmetric LOCC measurement.

PACS numbers: 03.65.Wj, 03.65.Ud, 02.20.-a

I. INTRODUCTION

Testing of maximally entangled state is a useful method for guaranteeing the quality of generated maximally entangled states. However, if we require a group symmetric condition for this method, the optimal test often requires infinite-valued measurement. Since any realizable measurement has a finite number of outcomes, it is needed to discretize the optimal measurement.

Now, we focus on the bipartite system $\mathcal{H}_d \otimes \mathcal{H}_d$, in which, the party A and B have the computational bases $\{|i\rangle_A\}_{i=0}^{d-1}$ and $\{|i\rangle_B\}_{i=0}^{d-1}$, respectively. When our target is testing whether the generated state is sufficiently close to the maximal entangled state

$$|\phi_{AB}^0\rangle = \frac{1}{\sqrt{d}} \sum_{i=0}^{d-1} |i\rangle_A \otimes |i\rangle_B$$

under a group symmetric condition, the optimal test can be given by

$$\begin{aligned} T_{inv}^{1,A \rightarrow B} &:= \int d|\varphi\rangle\langle\varphi| \nu(d\varphi) \\ &= |\phi_{A,B}^0\rangle\langle\phi_{A,B}^0| + \frac{1}{d+1}(I - |\phi_{A,B}^0\rangle\langle\phi_{A,B}^0|), \end{aligned} \quad (1)$$

where ν is the group invariant probability measure on the set of pure states, and φ and $\bar{\varphi}$ are given as $\varphi = \sum_{i=0}^{d-1} \varphi_i |i\rangle_A$ and $\bar{\varphi} = \sum_{i=0}^{d-1} \bar{\varphi}_i |i\rangle_B$. This measurement can be realized by the following procedure. In the first step, the system A performs the local group covariant measurement $\int d|\varphi\rangle\langle\varphi| \nu(d\varphi)$, and sends the system B the outcome φ . In the second step, the system B performs the two-valued measurement $\{|\bar{\varphi}\rangle\langle\bar{\varphi}|, I - |\bar{\varphi}\rangle\langle\bar{\varphi}|\}$. When Bob obtains the event corresponding to $\{|\bar{\varphi}\rangle\langle\bar{\varphi}|\}$, we support the maximal entangled state $|\phi_{AB}^0\rangle$.

This detection procedure can be generalize as follows. First, Alice performs a local measurement:

$$M = \{p_i |u_i\rangle\langle u_i|\}_i, \quad \|u_i\| = 1, \quad 0 \leq p_i \leq 1$$

and sends Bob the outcome i . Bob perform the two-valued measurement $\{|\bar{u}_i\rangle\langle\bar{u}_i|, I - |\bar{u}_i\rangle\langle\bar{u}_i|\}$. When Bob

obtain the event corresponding to $|\bar{u}_i\rangle\langle\bar{u}_i|$, we support the maximal entangled state $|\phi_{AB}^0\rangle$. This test can be written as the positive semi-definite matrix $T(M)$:

$$T(M) \stackrel{\text{def}}{=} \sum_i p_i |u_i \otimes \bar{u}_i\rangle\langle u_i \otimes \bar{u}_i|. \quad (2)$$

Indeed, when the local dimension d is 2, D'Ariano et al. [1] and Hayashi et al. [2] obtained the discrete own-way LOCC realization of the test $T_{inv}^{1,A \rightarrow B}$ as test $T(M)$ with an appropriate choice of the local measurement M . However, its general dimensional case was an open problem. In this paper, employing the concepts of symmetric informationally complete POVM (SIC-POVM) and mutually unbiased bases (MUB), we propose discrete own-way LOCC realizations of $T_{inv}^{1,A \rightarrow B}$. Also, the optimality of the proposed realization scheme is shown.

Next, we consider the case when Alice's system (Bob's system) is given as $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2}$ ($\mathcal{H}_{B_1} \otimes \mathcal{H}_{B_2}$) and the dimensions of all components coincide, i.e., $\dim \mathcal{H}_{A_1} = \dim \mathcal{H}_{A_2} = \dim \mathcal{H}_{B_1} = \dim \mathcal{H}_{B_2} = d$. In this case, we focus on the covariant POVM $M_{cov,u}^2$:

$$\begin{aligned} &M_{cov,u}^2(dg_1 dg_2) \\ &\stackrel{\text{def}}{=} d^2(g_1 \otimes g_2)|u\rangle\langle u|(g_1 \otimes g_2)^* \nu(dg_1) \nu(dg_2), \end{aligned}$$

where the vector u is a maximally entangled state and ν is the group invariant probability measure on $SU(d)$. The optimal test is given as the test $T_{inv}^{2,A \rightarrow B} \stackrel{\text{def}}{=} T(M_{cov,u}^2)$, which has the form [3]:

$$\begin{aligned} &T_{inv}^{2,A \rightarrow B} \\ &= |\phi_{A,B}^0\rangle\langle\phi_{A,B}^0| \otimes |\phi_{A,B}^0\rangle\langle\phi_{A,B}^0| \\ &\quad + \frac{1}{d^2 - 1}(I - |\phi_{A,B}^0\rangle\langle\phi_{A,B}^0|) \otimes (I - |\phi_{A,B}^0\rangle\langle\phi_{A,B}^0|). \end{aligned} \quad (3)$$

Indeed, The positive semi-definite matrix $T_{inv}^{2,A \rightarrow B}$ does not depend on the choice of the maximally entangled state u . In this paper, employing the concept of Clifford group, we provide a discrete own-way LOCC realization of $T_{inv}^{2,A \rightarrow B}$ when the local system is given as a composite system of a prime-dimensional system. Also, the optimality of the proposed realization scheme is shown.

*Electronic address: hayashi@math.is.tohoku.ac.jp

II. DISCRETE OWN-WAY LOCC REALIZATION OF $T_{inv}^{1,A \rightarrow B}$

A. Realizing scheme by SIC-POVM

In order to design the test $T_{inv}^{1,A \rightarrow B}$, we focus on the concept “symmetric informationally complete POVM (SIC-POVM)”. A rank-one POVM $\{p_i|u_i\rangle\langle u_i|\}$ on $\mathcal{H}_A = \mathbb{C}^d$ is called a *symmetric informationally complete POVM (SIC-POVM)*, if it satisfies the following conditions:

$$\begin{aligned} \#\{i\} &= d^2, \\ p_i &= \frac{1}{d} \\ |\langle u_i|u_j\rangle|^2 &= \frac{1}{d+1} \text{ for } i \neq j \end{aligned} \quad (4)$$

Currently, an SIC-POVM analytically is constructed when the dimension d is 2,3[5, 7],4[4, 7],5[7],6[6],7[8], 8[5], or 19[8]. Also, its existence is numerically verified up to $d = 45$ [4]. As is shown in Appendix A, any SIC-POVM $M_{sic} = \{p_i|u_i\rangle\langle u_i|\}_i$ satisfies

$$T(M_{sic}) = T_{inv}^{1,A \rightarrow B}, \quad (5)$$

that is, the test $T_{inv}^{1,A \rightarrow B}$ can be realized by an SIC-POVM. Moreover, if a POVM $M = \{M_i\}_i$ on $\mathcal{H}_A$ satisfies

$$T(M) = T_{inv}^{1,A \rightarrow B},$$

the inequality

$$\#\{i\} \geq d^2$$

holds. This is because the rank of the operator $T_{inv}^{1,A \rightarrow B}$ (which equal d^2) is less than the number of the elements of POVM M_i . Hence, we obtain

$$\min\{\#\{i\} | T(\{M_i\}_i) = T_{inv}^{1,A \rightarrow B}\} = d^2 \quad (6)$$

if there exists an SIC-POVM on $\mathbb{C}^d$. That is, the proposed realizing scheme by SIC-POVM is optimal in the sense of (6).

B. Realizing scheme by MUB

However, any SIC-POVM is not a randomized combination of projection valued measures as well as a projection valued measure. Since a projection valued measure (PVM) are more realizable than other POVM, it is more desired to design Alice’s POVM as a randomized combination of PVMs. For this purpose, we focus on mutually unbiased bases. $d+1$ orthonormal bases $\{\mathcal{B}_1, \dots, \mathcal{B}_{d+1}\}$ are called mutually unbiased bases (MUB) if

$$|\langle u|v\rangle|^2 = \frac{1}{d}, \forall u \in \mathcal{B}_i, \forall v \in \mathcal{B}_j, i \neq j.$$

The existence of MUB is shown when d is a prime[9] or a prime power[10]. Bandyopadhyay *et al.* gave a more explicit form in these cases [11]. Any mutually unbiased bases $\{\mathcal{B}_1, \dots, \mathcal{B}_{d+1}\}$ make the POVM $M_{\mathcal{B}_1, \dots, \mathcal{B}_{d+1}}$, i.e.,

$$M_{\mathcal{B}_1, \dots, \mathcal{B}_{d+1}} = \left\{ \frac{1}{d+1} |u_{i,j}\rangle\langle u_{i,j}| \right\}_{i,j},$$

where $\mathcal{B}_j = \{u_{1,j}, \dots, u_{d,j}\}$. This POVM always produces the desired test $T_{inv}^{1,A \rightarrow B}$ as

$$T(M_{\mathcal{B}_1, \dots, \mathcal{B}_{d+1}}) = T_{inv}^{1,A \rightarrow B}, \quad (7)$$

which is shown in Appendix B. This construction of the test $T_{inv}^{1,A \rightarrow B}$ is optimal in the following sense. Let $\{M^j\}$ be the set of projection-valued measures. A randomized combination of $\{M^j\}$, i.e., $M = \sum_j p_j M_j$ satisfies

$$T(M) = T_{inv}^{1,A \rightarrow B}. \quad (8)$$

Then, as is proven in Appendix C,

$$\#\{j\} \geq d+1, \quad (9)$$

which implies the optimality of the POVM consisting of MUB. Hence,

$$\min_{M_j: \text{PVM}} \left\{ \#\{j\} \mid T\left(\sum p_j M_j\right) = T_{inv}^{1,A \rightarrow B} \right\} = d+1 \quad (10)$$

if d is a prime or a prime power. That is, the proposed realizing scheme by MUB is optimal in the sense of (10).

III. DISCRETE OWN-WAY LOCC REALIZATION OF $T_{inv}^{2,A \rightarrow B}$

Next, we proceed to the case when both local systems consist of two subsystems. Given a finite group G and its projective representation f on $\mathcal{H}_{A_1} = \mathbb{C}^d$, by regarding $\mathcal{H}_{A_2}$ as the dual space of $\mathcal{H}_{A_1}$, the matrix $f(g)$ can be regarded as an element $|f(g)\rangle$ of $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2}$.

Theorem 1 *We assume the two conditions: (1) The representation f is irreducible. (2) The action $f \otimes \bar{f}$ of G to $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2}$ has only two irreducible components, i.e., the irreducible subspaces of $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2}$ for the action*

$$v_1 \otimes v_2 \rightarrow f(g)v_1 \otimes \overline{f(g)}v_2$$

are only the one-dimensional space $\langle \phi_{A_1, A_2}^0 \rangle$ and its orthogonal space $\langle \phi_{A_1, A_2}^0 \rangle^\perp$. Then, the resolution $M_f = \left\{ \frac{d^2}{|G|} \left| \frac{1}{\sqrt{d}} f(g) \right\rangle \left\langle \frac{1}{\sqrt{d}} f(g) \right| \right\}_{g \in G}$ satisfies the condition for a POVM, and

$$T(M_f) = T_{inv}^{2,A \rightarrow B}. \quad (11)$$

Its proof is given in Appendix D. This theorem yields a discrete own-way LOCC realization of $T_{inv}^{2,A \rightarrow B}$ from the representation f satisfying the above two conditions.

For example, Clifford group satisfies this assumption. For readers' convenience, we give its definition and prove that Clifford group satisfies this assumption. Clifford group $C(d)$ for d -dimensional system is given by

$$\begin{aligned} C(d) &:= \{U \in \mathcal{U}(d) | U \text{GP}(d)U^\dagger = \text{GP}(d)\} \\ \text{GP}(d) &:= \{e^{\sqrt{-1}\xi} W(i, j) | \xi \in \mathbb{R}, i, j \in \mathbb{Z}\} \\ I(d) &:= \{e^{\sqrt{-1}\xi} | \xi \in \mathbb{R}\}, \end{aligned}$$

where

$$\begin{aligned} Z &:= \sum_{j=0}^{d-1} \omega^j |j\rangle\langle j|, \quad X := \sum_{j=0}^{d-1} |j+1\rangle\langle j| \\ W(i, j) &:= X^i Z^j \end{aligned}$$

and ω is the d -th root of 1. As is shown in Appendix E, the natural representation of the group $C(d)$ satisfies the conditions (1) and (2). Then, the natural projective representation of the group $C(d)/I(d)$ also satisfies the conditions (1) and (2). As is shown in Lemma 5 in Appleby [8], when d is prime, the cardinality $|C(d)/I(d)|$ is $d^3(d^2 - 1)$. In the general case,

$$|C(d)/I(d)| = d^2 \left(\sum_{n=0}^{d-1} \nu(n, d) \nu(n+1, d) \right),$$

where $\nu(n, d)$ is the number of distinct ordered pairs $(x, y) \in \mathbb{Z}_d^2$ such that $xy = n \pmod{d}$.

IV. DISCUSSION

This paper has treated discretization of onw-way LOCC protocols. Using the concepts of symmetric informationally complete POVM (SIC-POVM), mutually unbiased bases (MUB), and Clifford group, we have proposed discrete own-way LOCC realizations of $T_{inv}^{1,A \rightarrow B}$ and $T_{inv}^{2,A \rightarrow B}$. This result indicates the importance of these concept in discrete mathematics. Since the existence of SIC-POVM and MUB is proven in limited cases, we cannot construct a discrete own-way LOCC realization of $T_{inv}^{1,A \rightarrow B}$ in the general case. Thus, further investigation for these concepts are required.

While the optimal test is given as $T_{inv}^{3,A \rightarrow B}$ when the local system consists of three subsystems by Hayashi [3], its discretization has not been obtained. Since the optimal test $T_{inv}^{3,A \rightarrow B}$ is closely related to GHZ state[3], its discretization may be related to GHZ state. Its construction remains as a future research.

Further, the optimal protocol is often given as a protocol with infinite elements in quantum information. In such a case, it is required to discretize this protocol. This kind of discretization is an interesting interdisciplinary topic between quantum information and discrete mathematics.

Acknowledgment

This research was partially supported by a Grant-in-Aid for Scientific Research on Priority Area 'Deepening and Expansion of Statistical Mechanical Informatics (DEX-SMI)', No. 18079014 and a MEXT Grant-in-Aid for Young Scientists (A) No. 20686026.

APPENDIX A: PROOF OF (5)

First, we show that $u_1 \otimes \overline{u_1}, \dots, u_{d^2} \otimes \overline{u_{d^2}}$ are linearly independent. We choose complex numbers $a_1, \dots, a_{d^2}$ such that

$$\sum_i a_i u_i \otimes \overline{u_i} = 0.$$

Taking trace, we have

$$a_1 + \sum_{i \neq 1} a_i = 0.$$

On the other hand,

$$0 = \langle u_1 \otimes \overline{u_1} | \sum_i a_i u_i \otimes \overline{u_i} \rangle = a_1 + \frac{1}{d+1} \sum_{i \neq 1} a_i.$$

Hence, we obtain $a_1 = 0$. Similarly, we can show $a_i = 0$, which implies the linear independence.

Since the dimension of $\mathcal{H}_A \otimes \mathcal{H}_B$ is d^2 , any element of $\mathcal{H}_A \otimes \mathcal{H}_B$ can be expressed as

$$\sum_j a_j u_j \otimes \overline{u_j}.$$

We can calculate

$$\begin{aligned} & \left\langle \sum_i a_i u_i \otimes \overline{u_i} \middle| T(M_{sic}) \middle| \sum_j a_j u_j \otimes \overline{u_j} \right\rangle \\ &= \left\langle \sum_i a_i u_i \otimes \overline{u_i} \middle| \left(\sum_k \frac{1}{d} |u_k \otimes \overline{u_k}\rangle \langle u_k \otimes \overline{u_k}| \right) \middle| \sum_j a_j u_j \otimes \overline{u_j} \right\rangle \\ &= \frac{d+2}{(d+1)^2} \left| \sum_k a_k \right|^2 + \frac{d}{(d+1)^2} \sum_k |a_k|^2. \end{aligned}$$

On the other hand, its norm is calculated as

$$\left\| \sum_j a_j u_j \otimes \overline{u_j} \right\|^2 = \frac{1}{d+1} \left| \sum_k a_k \right|^2 + \frac{d}{d+1} \sum_k |a_k|^2.$$

Since

$$\left| \langle \phi_{A,B}^0 | \sum_j a_j u_j \otimes \overline{u_j} \rangle \right|^2 = \frac{1}{d} \left| \sum_j a_j \right|^2,$$

we obtain

$$\begin{aligned}
& \left\langle \sum_i a_i u_i \otimes \overline{u_i} \left| T_{inv}^{1,A \rightarrow B} \right| \sum_j a_j u_j \otimes \overline{u_j} \right\rangle \\
&= \left\langle \sum_i a_i u_i \otimes \overline{u_i} \left| \left(\frac{d}{d+1} |\phi_{A,B}^0\rangle \langle \phi_{A,B}^0| + \frac{1}{d+1} I \right) \right. \right. \\
&\quad \left. \left. \left| \sum_j a_j u_j \otimes \overline{u_j} \right\rangle \right\rangle \\
&= \frac{d}{d+1} \frac{1}{d} \left| \sum_j a_j \right|^2 \\
&\quad + \frac{1}{d+1} \left(\frac{1}{d+1} \left| \sum_k a_k \right|^2 + \frac{d}{d+1} \sum_k |a_k|^2 \right) \\
&= \left\langle \sum_i a_i u_i \otimes \overline{u_i} \left| T(M_{sic}) \right| \sum_j a_j u_j \otimes \overline{u_j} \right\rangle.
\end{aligned}$$

Therefore, we obtain (5).

APPENDIX B: PROOF OF (7)

We focus on the subspace $\langle \phi_{A,B}^0 \rangle^\perp$ orthogonal to $\phi_{A,B}^0$. The subspace $\mathcal{B}'_j = \langle u_{1,j} \otimes \overline{u_{1,j}} - \frac{1}{d} \phi_{A,B}^0, \dots, u_{d-1,j} \otimes \overline{u_{d-1,j}} - \frac{1}{d} \phi_{A,B}^0 \rangle$ belongs to the subspace $\langle \phi_{A,B}^0 \rangle^\perp$, and its dimension is $d-1$. Since

$$\langle u_{i,j} \otimes \overline{u_{i,j}} - \frac{1}{d} \phi_{A,B}^0 | u_{i',j'} \otimes \overline{u_{i',j'}} - \frac{1}{d} \phi_{A,B}^0 \rangle = 0, \quad j \neq j', \quad (\text{B1})$$

The spaces $\mathcal{B}'_1, \dots, \mathcal{B}'_{d+1}$ are orthogonal to each other. Since the dimension of the subspace $\langle \phi_{A,B}^0 \rangle^\perp$ is $d^2 - 1$, the subspace $\langle \phi_{A,B}^0 \rangle^\perp$ is spanned by the spaces $\mathcal{B}'_1, \dots, \mathcal{B}'_{d+1}$. Therefore, any element of the space $\mathcal{H}_A \otimes \mathcal{H}_B$ can be expressed as $\sum_{j=1}^{d+1} \sum_{i=1}^d a_{i,j} u_{i,j} \otimes \overline{u_{i,j}}$. In the following, we abbreviate the sum $\sum_{j=1}^{d+1} \sum_{i=1}^d$ as $\sum_{j,i}$.

We calculate

$$\begin{aligned}
& \left\langle \sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \left| T(M_{\mathcal{B}_1, \dots, \mathcal{B}_{d+1}}) \right| \sum_{j',i'} a_{i',j'} u_{i',j'} \otimes \overline{u_{i',j'}} \right\rangle \\
&= \left\langle \sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \left| \left(\sum_{l,k} \frac{1}{d+1} |u_{k,l} \otimes \overline{u_{k,l}}\rangle \langle u_{k,l} \otimes \overline{u_{k,l}}| \right) \right. \right. \\
&\quad \left. \left. \left| \sum_{j',i'} a_{i',j'} u_{i',j'} \otimes \overline{u_{i',j'}} \right\rangle \right\rangle \\
&= \frac{1}{d+1} \sum_{l,k} \left| \sum_{j,i} \langle u_{k,l} \otimes \overline{u_{k,l}} | a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \rangle \right|^2 \\
&= \frac{1}{d} \left| \sum_{j,i} a_{i,j} \right|^2 - \frac{1}{d(d+1)} \sum_j \left| \sum_i a_{i,j} \right|^2 + \frac{1}{d+1} \sum_{j,i} |a_{i,j}|^2
\end{aligned}$$

On the other hand, its norm is calculated as

$$\begin{aligned}
& \left\| \sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \right\|^2 \\
&= \frac{1}{d} \left| \sum_{j,i} a_{i,j} \right|^2 - \frac{1}{d} \sum_j \left| \sum_i a_{i,j} \right|^2 + \sum_{j,i} |a_{i,j}|^2
\end{aligned}$$

Since

$$\left| \left\langle \phi_{A,B}^0 \left| \sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \right\rangle \right|^2 = \frac{1}{d} |a_{i,j} u_{i,j}|^2,$$

we obtain

$$\begin{aligned}
& \left\langle \sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \left| T_{inv}^{1,A \rightarrow B} \right| \sum_{j',i'} a_{i',j'} u_{i',j'} \otimes \overline{u_{i',j'}} \right\rangle \\
&= \left\langle \sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \left| \frac{d}{d+1} |\phi_{A,B}^0\rangle \langle \phi_{A,B}^0| + \frac{1}{d+1} I \right. \right. \\
&\quad \left. \left. \left| \sum_{j',i'} a_{i',j'} u_{i',j'} \otimes \overline{u_{i',j'}} \right\rangle \right\rangle \\
&= \frac{1}{d} \left| \sum_{j,i} a_{i,j} \right|^2 - \frac{1}{d(d+1)} \sum_j \left| \sum_i a_{i,j} \right|^2 + \frac{1}{d+1} \sum_{j,i} |a_{i,j}|^2 \\
&= \left\langle \sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \left| T(M_{\mathcal{B}_1, \dots, \mathcal{B}_{d+1}}) \right| \sum_{j',i'} a_{i',j'} u_{i',j'} \otimes \overline{u_{i',j'}} \right\rangle.
\end{aligned}$$

Therefore, we obtain (7).

APPENDIX C: PROOF OF (9)

Let $M^j = \{|u_{i,j}\rangle\langle u_{i,j}|\}$. We focus on the projection P to the subspace $\langle \phi_{A,B}^0 \rangle^\perp$ orthogonal to $\phi_{A,B}^0$ and the subspace $\mathcal{B}_j'' \stackrel{\text{def}}{=} \langle u_{1,j} \otimes \overline{u_{1,j}}, \dots, u_{d,j} \otimes \overline{u_{d,j}} \rangle$. The image $P\mathcal{B}_j''$ is $\langle u_{1,j} \otimes \overline{u_{1,j}} - \frac{1}{d}\phi_{A,B}^0, \dots, u_{d-1,j} \otimes \overline{u_{d-1,j}} - \frac{1}{d}\phi_{A,B}^0 \rangle$. The condition (8) implies that the sum of the rank of the space $P\mathcal{B}_j''$ is greater than $d^2 - 1$, *i.e.*, the dimension of the space $\langle \phi_{A,B}^0 \rangle^\perp$. Thus, $\#\{j\}(d-1) \geq d^2 - 1$, which implies the inequality (9).

APPENDIX D: PROOF OF THEOREM 1

First, we prove that M_f satisfies the condition for POVM. The irreducibility of the action f guarantees that

$$\begin{aligned} & \frac{d}{|G|} \sum_{g \in G} \langle k|f(g)|l\rangle \langle l'|f(g)|k'\rangle \\ &= \langle k| \left(\frac{d}{|G|} \sum_{g \in G} f(g)|l\rangle \langle l'|f(g) \right) |k'\rangle \\ &= \langle k|\langle l|l'\rangle I|k'\rangle = \delta_{k,k'} \delta_{l,l'}. \end{aligned}$$

we obtain

$$\begin{aligned} & \frac{d}{|G|} \sum_{g \in G} \left| \langle f(g)| \left(\sum_{k,l} a_{k,l} |k\rangle \otimes |l\rangle \right) \right|^2 \\ &= \frac{d}{|G|} \sum_{g \in G} \sum_{k,l} \sum_{k',l'} a_{k,l} \overline{a_{k',l'}} \langle k|f(g)|l\rangle \langle l'|f(g)|k'\rangle \\ &= \sum_{k,l} a_{k,l} \overline{a_{k,l}}, \end{aligned}$$

which implies

$$\frac{d^2}{|G|} \sum_{g \in G} \left| \left\langle \frac{1}{\sqrt{d}} f(g) \right\rangle \left\langle \frac{1}{\sqrt{d}} f(g) \right\rangle \right| = I_{A_1, A_2}.$$

Hence, $M_f = \left\{ \frac{d^2}{|G|} \left| \left\langle \frac{1}{\sqrt{d}} f(g) \right\rangle \left\langle \frac{1}{\sqrt{d}} f(g) \right\rangle \right| \right\}_{g \in G}$ is a POVM.

Next, we show (11). We focus on the action of the group $G \times G$ to the total space $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2} \otimes \mathcal{H}_{B_1} \otimes \mathcal{H}_{B_2}$ as

$$\begin{aligned} & u_1 \otimes u_2 \otimes v_1 \otimes v_2 \\ & \mapsto f(g_1)u_1 \otimes \overline{f(g_2)u_2} \otimes \overline{f(g_1)v_1} \otimes f(g_2)v_2 \end{aligned}$$

for $u_i \in \mathcal{H}_{A_i}$, $v_i \in \mathcal{H}_{B_i}$, and any pair $(g_1, g_2) \in G \times G$. Due to the condition (2), the irreducible decomposition of the space $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2} \otimes \mathcal{H}_{B_1} \otimes \mathcal{H}_{B_2}$ is given as $\langle \phi_{A_1, B_1}^0 \rangle \otimes \langle \phi_{A_2, B_2}^0 \rangle \oplus \langle \phi_{A_1, B_1}^0 \rangle \otimes \langle \phi_{A_2, B_2}^0 \rangle^\perp \oplus \langle \phi_{A_1, B_1}^0 \rangle^\perp \otimes \langle \phi_{A_2, B_2}^0 \rangle \oplus \langle \phi_{A_1, B_1}^0 \rangle^\perp \otimes \langle \phi_{A_2, B_2}^0 \rangle^\perp$.

As is checked below, the test $T(M_f)$ is invariant for this action:

$$\begin{aligned} & f(g_1) \otimes \overline{f(g_2)} \otimes \overline{f(g_1)} \otimes f(g_2) T(M_f) \left(f(g_1) \otimes \overline{f(g_2)} \otimes \overline{f(g_1)} \otimes f(g_2) \right)^\dagger \\ &= \frac{d^2}{|G|} \sum_{g \in G} \left| \frac{1}{d} f(g_1) f(g) f(g_2)^\dagger \otimes \overline{f(g_1) f(g) f(g_2)^\dagger} \right\rangle \left\langle \frac{1}{d} f(g_1) f(g) f(g_2)^\dagger \otimes \overline{f(g_1) f(g) f(g_2)^\dagger} \right| \\ &= \frac{d^2}{|G|} \sum_{g' \in G} \left| \frac{1}{d} f(g') \otimes \overline{f(g')} \right\rangle \left\langle \frac{1}{d} f(g') \otimes \overline{f(g')} \right| = T(M_f), \end{aligned}$$

where we denote $g_1 g_2^{-1}$ by g' . Hence, the test $T(M_f)$

has the form

$$\begin{aligned} & T(M_f) \\ &= a |\phi_{A_1, B_1}^0\rangle \langle \phi_{A_1, B_1}^0| \otimes |\phi_{A_2, B_2}^0\rangle \langle \phi_{A_2, B_2}^0| \\ & \quad + b (I - |\phi_{A_1, B_1}^0\rangle \langle \phi_{A_1, B_1}^0|) \otimes |\phi_{A_2, B_2}^0\rangle \langle \phi_{A_2, B_2}^0| \\ & \quad + c |\phi_{A_1, B_1}^0\rangle \langle \phi_{A_1, B_1}^0| \otimes (I - |\phi_{A_2, B_2}^0\rangle \langle \phi_{A_2, B_2}^0|) \\ & \quad + d (I - |\phi_{A_1, B_1}^0\rangle \langle \phi_{A_1, B_1}^0|) \otimes (I - |\phi_{A_2, B_2}^0\rangle \langle \phi_{A_2, B_2}^0|). \end{aligned}$$

Since $f(g)$ is the unitary matrix, $\frac{1}{\sqrt{d}}f(g)$ is a maximally entangled state on $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2}$. Since $\left|\frac{1}{\sqrt{d}}f(g)\right\rangle$ is maximally entangled, Lemma 5 in Hayashi [3] yields that

$$T(M_f) = |\phi_{A_1, B_1}^0 \otimes \phi_{A_2, B_2}^0\rangle\langle\phi_{A_1, B_1}^0 \otimes \phi_{A_2, B_2}^0| + PT(M_f)P, \quad (\text{D1})$$

where

$$P \stackrel{\text{def}}{=} (I - |\phi_{A_2, B_2}^0\rangle\langle\phi_{A_2, B_2}^0|) \otimes (I - |\phi_{A_1, B_1}^0\rangle\langle\phi_{A_1, B_1}^0|).$$

This relation implies that $b = c = 0$. Thus, the relation $\text{Tr } T(M_f) = d^2$ yields

$$\begin{aligned} T(M_f) &= |\phi_{A_1, B_1}^0\rangle\langle\phi_{A_1, B_1}^0| \otimes |\phi_{A_2, B_2}^0\rangle\langle\phi_{A_2, B_2}^0| \\ &\quad + \frac{1}{d^2 - 1} (I - |\phi_{A_1, B_1}^0\rangle\langle\phi_{A_1, B_1}^0|) \otimes (I - |\phi_{A_2, B_2}^0\rangle\langle\phi_{A_2, B_2}^0|), \end{aligned}$$

which implies (11).

APPENDIX E: PROOF OF IRREDUCIBILITY

It is known that the natural representation of the subgroup $\text{GP}(d) \subset \text{C}(d)$ satisfies the condition (1). Hence,

it is sufficient to show the condition (2). the irreducible spaces of the subgroup $\text{GP}(d) \subset \text{C}(d)$ are d^2 one-dimensional subspaces generated by $|W(i, j)\rangle$ for i, j . The representation of $\text{GP}(d)$ on each irreducible subspaces is different. Thus, the irreducible subspace of the larger group $\text{C}(d)$ should be represented as the direct sum of these subspaces. As is shown in Lemma 1 in Appleby [8], for any (i, j) and any $F \in \text{SL}(2, \mathbb{Z}_{\bar{d}})$, there exists an element $U \in \text{C}(d)$ such that $f(U) \otimes \overline{f(U)}|W(i, j)\rangle = e^{\sqrt{-1}\delta_{i,j}, F}|W(F(i, j))\rangle$, where

$$\bar{d} := \begin{cases} d & \text{if } d \text{ is odd} \\ 2d & \text{if } d \text{ is even.} \end{cases} \quad (\text{E1})$$

For any pair $(i, j) \neq (0, 0)$, there exists an element $F \in \text{SL}(2, \mathbb{Z}_{\bar{d}})$ such that $(i, j) = F(1, 0)$. Since any irreducible subspace should be spanned by the subset of $\{|W(i, j)\rangle\}_{i,j}$, the space spanned by $\{|W(i, j)\rangle\}_{(i,j) \neq (0,0)}$ is irreducible. Thus, the condition (2) holds.

-
- [1] G. M. D'Ariano, C. Macchiavello, and M. G. A. Paris, "Local observables for entanglement witnesses," *Phys. Rev. A* **67** 042310, 2003.
 - [2] M. Hayashi, K. Matsumoto, and Y. Tsuda, "A study of LOCC-detection of a maximally entangled state using hypothesis testing," *J. Phys. A: Math. and Gen.*, **39** 14427-14446 (2006).
 - [3] M. Hayashi, Group theoretical study of LOCC-detection of maximally entangled state using hypothesis testing, arXiv:0810.3380.
 - [4] J. M. Renes, R. Blume-Kohout, A. J. Scott, and C. M. Caves, "Symmetric Informationally Complete Quantum Measurements," *J. Math. Phys.*, **45**, 2171-2180 (2004); quant-ph/0310075.
 - [5] A. Koldobsky and H. König, "Aspects of the Isometric Theory of Banach Spaces," in *Handbook of Geometry of Banach Spaces*, Vol. 1, edited by W. B. Johnson and J. Lindenstrauss (North-Holland, Dordrecht, 2001), pp.899-939.
 - [6] Grassl, M., "On SIC-POVMs and MUBs in dimension 6," *Proceedings of EQIS'04*, pp. 60-61, (2004); quant-ph/0406175.
 - [7] Zauner, G., "Quantum designs—foundations of a non-commutative theory of designs," (in German), Ph.D. thesis, University of Vienna, (1999).
 - [8] Appleby, D M, "SIC-POVMs and the Extended Clifford Group," *J. Math. Phys.*, **46**, 052107 (2005).
 - [9] I. D. Ivanovic, "Geometrical description of quantum state determination," *J. Phys. A: Math. and Gen.*, **14**, No. 12, 3241-3245, (1981).
 - [10] W. K. Wootters and B. D. Fields, "Optimal state-determination by mutually unbiased measurements," *Annals of Physics*, **191**, No. 2, 363-381, (1989).
 - [11] S. Bandyopadhyay, P. O. Boykin, V. Roychowdhury, and F. Vatan, "A New Proof for the Existence of Mutually Unbiased Bases," *Algorithmica*, **34** (2002), pp. 512-528; quant-ph/0103162.